

Lenticulostriate vasculopathy in very preterm infants

L M Leijser, S J Steggerda, F T de Bruïne, et al.

Arch Dis Child Fetal Neonatal Ed 2010 95: F42-F46 originally published online May 19, 2009 doi: 10.1136/adc.2009.161935

Updated information and services can be found at: http://fn.bmj.com/content/95/1/F42.full.html

These include:

References	This article cites 32 articles, 6 of which can be accessed free at: http://fn.bmj.com/content/95/1/F42.full.html#ref-list-1
Email alerting service	Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To order reprints of this article go to: http://fn.bmj.com/cgi/reprintform ¹ Department of Pediatrics, Leiden University Medical Center, Leiden, The Netherlands; ² Department of Radiology, Leiden University Medical Center, Leiden, The Netherlands

Correspondence to: Lara M Leijser, Department of Pediatrics, Division of Neonatology, J6-S, Leiden University Medical Center, P.O. Box 9600, 2300 RC Leiden, The Netherlands; I.m.leijser@lumc.nl

Accepted 8 May 2009 Published Online First 19 May 2009 Downloaded from fn.bmj.com on May 4, 2010 - Published by group.bmj.com

Lenticulostriate vasculopathy in very preterm infants

L M Leijser,¹ S J Steggerda,¹ F T de Bruïne,² A van Zuijlen,¹ A van Steenis,¹ F J Walther,¹ G van Wezel-Meijler¹

ABSTRACT

Objective: To assess for lenticulostriate vasculopathy (LSV) on cranial ultrasound (cUS) scans of very preterm infants: incidence and aetiology, evolution during neonatal period, association with clinical parameters, and MRI equivalent. **Design:** Prospective study.

Setting: Tertiary neonatal referral centre.

Patients: Very preterm infants (<32 weeks) underwent sequential cUS throughout the neonatal period and MRI around term age. cUS were evaluated for LSV and other changes, and MRI for changes in signal and myelination in deep grey matter. LSV was divided into early-onset (≤7 postnatal days) and late-onset (>7 postnatal days). Perinatal clinical parameters were collected for all infants and compared between groups.

Results: In 22/111 (20%) infants LSV was detected: earlyonset in 5 and late-onset in 17. LSV mostly presented some weeks after birth and persisted for several months. There were no associations between LSV and other changes on cUS or deep grey matter changes on MRI. Infants with late-onset LSV were younger and smaller at birth than infants with early-onset LSV. Postmenstrual age at first detection was comparable for both LSV groups. There were no associations between LSV and perinatal clinical parameters, but infants with LSV had less episodes of hypotension than infants without LSV. **Conclusions:** LSV is a frequent finding on cUS in verv preterm infants, but does not show on MRI. The postmenstrual age, rather than gestational and postnatal age, seems important in LSV development. LSV is not associated with clinical parameters. When encountered in otherwise healthy preterm infants, LSV is probably a

Lenticulostriate vasculopathy (LSV), presenting as hyperechogenic vessels in the basal ganglia and/or thalamus (BGT) region, can be detected on cranial ultrasound (cUS) scans of neonates.¹ The incidence of LSV in neonates varies between 0.3 and 32%.²⁻¹⁶ LSV can be unilateral or bilateral, and punctate, linear or branching.^{4 5 9 10 17} Histological studies show intramural and perivascular depositions of amorphous basophilic material in the lenticulostriate vessels, probably causing the hyperechogenic appearance on cUS. The vessels are usually medium sized, having thickened and hypercellular walls without fibrosis or hyalinisation.^{2 5 9 13}

benign temporary phenomenon.

LSV has been associated with several clinical conditions, including infections, both congenital (eg, Toxoplasmosis, Other, Rubella, Cytomegalovirus, Herpes (including Parvo B19) (TORCH) infections) and acquired, chromosomal abnormalities, congenital heart disease, other congenital malformations, hypoxic-ischaemic events, fetal alcohol and drug exposure, twin-to-twin transfusion syndrome, neonatal lupus erythematosus, hydrops fetalis, metabolic disorders and diabetic fetopathy.²⁻¹⁰ ¹² ¹³ ¹⁵ ¹⁷⁻²⁸

What is already known on this topic

- Lenticulostriate vasculopathy (LSV), presenting as hyperechogenic vessels in the deep grey matter, can be detected on cranial ultrasound scans of neonates and its incidence varies between 0.3 and 32%.
- LSV has been associated with several clinical conditions of the fetus and neonate, but its aetiology and clinical significance remain largely unclear.
- Studies on the incidence, evolution and aetiology of LSV and associated clinical parameters in preterm neonates are limited.

What this study adds

- In very preterm infants, lenticulostriate vasculopathy (LSV) is a frequent cranial ultrasound finding, mostly presenting some weeks after birth and persisting for several months.
- Postmenstrual age, rather than gestational and postnatal age, seems important in LSV development; LSV is not associated with perinatal clinical parameters, including congenital infections.
- In healthy very preterm infants, LSV may be a benign temporary phenomenon.

It occurs more often in infants of multiple than of singleton pregnancies and in full-term than in preterm neonates.^{12 13 25 26} However, the aetiology and clinical significance of LSV are largely unclear.^{8 13 15}

Prospective studies on the incidence and aetiology of LSV and associated clinical parameters in preterm infants are limited.^{4 5 10 12 14 15} The aims of our prospective study were to assess the incidence, aetiology and evolution of LSV in a large cohort of very preterm infants. Additional aims were to assess its association with various perinatal clinical parameters, previously associated with brain injury in preterm infants, presence of an equivalent on MRI, and presence of changes in maturation of the BGT in infants with as compared to without LSV.

PATIENTS AND METHODS Patients

Very preterm infants (gestational age (GA) <32 weeks), admitted to the tertiary neonatal unit of the Leiden University Medical Center between May 2006 and October 2007, were eligible for a neuroimaging study, assessing and comparing brain imaging findings on cUS and MRI. For this part of the study, emphasis was placed on LSV.

Exclusion criteria were congenital anomalies of the central nervous system, severe other congenital anomalies, chromosomal and metabolic disorders, and neonatal meningitis.

Cranial ultrasound

As part of the routine care, sequential cUS scans were performed by the attending (fellow) neonatologists, using an Aloka $\alpha 10$ scanner with multifrequency transducer (Biomedic Nederland B.V., Almere, the Netherlands), within 24 h of birth, at least weekly from the day of birth or admission until discharge or transfer to another hospital, and on the day of the term equivalent MRI examination.²⁹ Of included infants, all cUS scans were assessed as recently described.³⁰

LSV was considered present when on at least one cUS examination, a punctate or linear echogenic structure was seen in the distribution of the thalamo-striatal vessels in the BGT region, on both coronal and (para)sagittal views (fig 1). The side (left and/or right), appearance (punctate, linear and/or branching), and location (basal ganglia and/or thalamus) were recorded. The day LSV was first detected on sequential cUS and its evolution and duration (up to the day of the term equivalent cUS) were noted. As previous studies described differences between preterm infants with LSV presenting within the first 7–10 days of birth and thereafter,¹¹ we divided LSV into early-onset (first seen on initial or subsequent cUS performed \leq 7 days of birth) and late-onset (first seen on cUS performed >7 days of birth).

Other changes, including echodensities in the frontal white matter (frontal echodensities), periventricular echodensities, cystic white matter lesions, intraventricular haemorrhage, post-haemorrhagic ventricular dilatation, and periventricular haemorrhagic infarction, were recorded.³⁰

Magnetic resonance imaging

MRI examinations were performed in all very preterm infants, preferably around or just after term equivalent age (TEA), according to our standard protocol, using a 3.0 Tesla Philips MR system (Philips Medical Systems, Best, the Netherlands) as recently described.³¹ For infants who were still unstable and/or ventilator dependent around that age, MRI was postponed. For this part of the neuroimaging study, only the T₁-, T₂-, and susceptibility-weighted sequences were analysed.

All MRIs were assessed as recently described.³⁰ Special attention was paid to signal intensity changes in the BGT. The side (left and/or right), appearance, and location (basal ganglia and/or thalamus) were recorded. Myelination in the BGT region was compared to reference images, obtained with lower field strengths.^{32,35}

Clinical parameters

For all included infants the following perinatal clinical parameters were retrospectively collected: antenatal corticosteroid treatment; GA at birth; birth weight; gender; plurity, and in case of monochorionic twins presence of twin-to-twin transfusion syndrome; congenital TORCH infection based on positive maternal serum IgM or PCR screening; respiratory distress syndrome requiring mechanical ventilation and surfactant treatment; bronchopulmonary dysplasia, if oxygen dependence \geq 30% at 36 weeks' postmenstrual age (PMA)³⁶; dexamethasone treatment for bronchopulmonary dysplasia; patent ductus arteriosus requiring medical and/or surgical treatment; hypotension requiring inotropic treatment; sepsis, if positive blood culture, and/or necrotising enterocolitis, if grade \geq 2a according to Bell *et al.*³⁷

Data analysis

Statistical analyses were performed using SPSS software V.14.0 (SPSS inc., Chicago, Illinois, USA). For the analyses, only infants who were admitted to our unit for at least 7 days were included. The incidence of LSV was calculated. Incidences of (other) changes on cUS and MRI were compared between infants with and without LSV, using a Pearson χ^2 test. Subsequently, infants with LSV were divided into early-onset and late-onset groups. Characteristics of LSV and incidences of (other) changes on cUS and MRI were compared between both LSV groups, using a Pearson χ^2 test.

Incidences of clinical parameters were compared between infants with and without LSV and between the early-onset and late-onset groups, using a Pearson χ^2 test for categorical and an unpaired t test for continuous variables. Clinical variables with p < 0.1 were entered into a stepwise logistic regression model to identify independent predictors of LSV. As several of the collected clinical parameters may be strongly related to GA at birth, this was always included in the model. Level of significance was $p \leq 0.05$.

Ethical approval

The study was approved by the Medical Ethics Committee and informed consent was obtained from the parents.

RESULTS

Patients

During the study period, 182 very preterm infants were eligible for the LSV study, of whom 130 (80 male) were included. Fiftytwo infants were excluded; three because of structural brain abnormalities and 49 because informed parental consent was not obtained. Reasons for not obtaining consent included transfer to another hospital or death within a very short period

Figure 1 Coronal (A) and sagittal (B) cranial ultrasound scans of a preterm infant (gestational age 26.4 weeks), scanned at postmenstrual age 30.7 weeks, showing bilateral lenticulostriate vasculopathy having a branching appearance on the right side and a linear appearance on the left side.



Original article

of birth, rejection of participation by the parents, and practical problems such as language barrier and travel distance to the hospital. Median GA and birth weight of included infants were 29.0 (range 25.6–31.9) weeks and 1141 (520–1960) g, respectively. There were no significant differences in GA and weight at birth between infants with and without informed consent.

In 111 (83.5%; 68 male) infants, sequential cUS (median number 8, range 4–21) were performed during a period of at least 7 days, but in 12 of these infants no or inadequate cUS and/or MRI around TEA were obtained. In three infants, MRI was performed elsewhere, using different protocols. In two infants, MR images were difficult to interpret due to movement artefacts, and in seven infants the parents withdrew participation from MRI. Therefore, in 99 infants (61 male) contemporaneous cUS and MRI were obtained at a median PMA of 43.4 (40.1–55.9) weeks. In 61 infants MRI was performed around TEA; in the other 38 between 44.0 and 55.9 weeks' PMA.

Cranial ultrasound (n = 111) LSV

LSV group

In 22 infants (19.8%; 11 male), LSV was detected on at least one cUS examination (median number 6, range 1–12). Median GA and birth weight were 27.9 (26.0–31.7) weeks and 1095 (620–1800) g, respectively. Characteristics of LSV are shown in table 1.

Early-onset and late-onset LSV

LSV was detected within the first 7 postnatal days in five infants (22.7%), while thereafter in 17 (77.3%). In three infants, LSV was seen on the first cUS scan performed within 24 h of birth. Characteristics of LSV and differences therein between infants with early- and late-onset LSV are shown in table 1. PMA at first detection of LSV was not different between both groups. In infants with early-onset LSV, LSV was significantly more often left-sided than in infants with late-onset LSV, in whom LSV was mostly right-sided. No other differences in characteristics were found.

Relation with other changes

In most very preterm infants other changes were detected on $cUS.^{so}$ No association was found between LSV, or first detection of LSV, and other changes on cUS.

MRI (n = 99)

In none of the infants with LSV, including infants with LSV seen on contemporaneous cUS, signal intensity changes were detected in corresponding areas on MRI. Myelination in the BGT region was normal in all infants as compared to reference images. No association was found between LSV and changes in signal or myelination in the BGT on MRI.

Relation between LSV and clinical parameters (n = 111) LSV group

Characteristics of perinatal clinical parameters for infants with and without LSV are shown in table 2. Infants with LSV had significantly less episodes of hypotension than infants without LSV. Hypotension was identified as an independent variable. No differences were found for the other clinical parameters.

Early-onset and late-onset LSV

Characteristics of clinical parameters for the early- and late-onset LSV groups are shown in table 2. Infants with late-onset LSV had a significantly lower GA and weight at birth than infants with early-onset LSV. Only GA was an independent variable. No differences were found for the other clinical parameters.

DISCUSSION

To our knowledge, this is the first prospective study describing the incidence, evolution and association with clinical parameters of LSV in a large cohort of very preterm infants.

Our incidence (20%) is higher than previously reported.^{3–9} ^{11–16} Chamnanvanakij *et al*⁸ found an incidence of 5.1% in a small group of preterm infants with birth weight <1250 g, and in a retrospective study in preterm infants <35 weeks' GA, an

 Table 1
 Characteristics of lenticulostriate vasculopathy (LSV) for the LSV group and for the early-onset and late-onset LSV groups separately

				Early-onset	
Characteristics of LSV	LSV group (n = 22)	Early-onset LSV group (n = 5)	Late-onset LSV group (n = 17)	versus late- onset p value	
Detection					
First, median (range)					
Pn age (days)	17.0 (0–117)	0.0 (0-7)	20.0 (13–117)	0.034	
PMA (weeks)	30.5 (27.9-44.1)	30.0 (27.9-31.4)	31.0 (28.1–44.1)	NS	
Last, median (range)					
Pn age (days)	97.0 (14–166)	62.0 (14-166)	106.0 (57–165)	NS	
PMA (weeks)	42.6 (31.9-55.1)	36.7 (31.9–55.1)	42.7 (35.0-50.4)	NS	
Duration (days), median (range)	69.5 (1-166)	62.0 (7-166)	71.0 (1–112)	NS	
On first cUS, n (%)	3 (13.6)	3 (60.0)	0 (0.0)	0.006	
On term cUS, n (%)	15 (68.2)	2 (40.0)	13 (76.5)	NS	
At first detection					
Laterality, n (%)					
Unilateral	15 (68.2)	4 (80.0)	11 (64.7)	NS	
Right	11 (73.3)	1 (25.0)	10 (90.9)	0.022	
Left	4 (26.7)	3 (75.0)	1 (9.1)		
Bilateral	7 (31.8)	1 (20.0)	6 (35.3)		
Appearance, n (%)					
Punctate	4 (18.2)	1 (20.0)	3 (17.6)	NS	
Linear	17 (77.3)	4 (80.0)	13 (76.5)		
Branching	1 (4.5)	0 (0)	1 (5.9)		

cUS, cranial ultrasound; NS, not significant; PMA, postmenstrual age; pn, postnatal.

	1 7 1			0 1			
Clinical parameters	Total group (n = 111)	LSV group (n = 22)	Non-LSV group (n = 89)	LSV versus non-LSV group p value	Early-onset LSV group (n = 5)	Late-onset LSV group (n = 17)	Early-versus late-onset p value
Antenatal corticosteroids, n (%)	66 (59.5)	16 (72.7)	50 (56.2)	NS	5 (100)	11 (64.7)	NS
GA (weeks), median (range)	28.7 (25.6–31.9)	27.9 (26.0–31.7)	29.0 (25.6–31.9)	NS	29.9 (27.9–31.4)	27.3 (26.0–31.7)	0.041*
Birth weight (g), median (range)	1072 (520–1960)	1095 (620–1800)	1065 (520-1960)	NS	1410 (1066–1750)	1030 (620–1800)	0.045
Male, n (%)	68 (61.3)	11 (50.0)	57 (64.0)	NS	2 (40.0)	9 (52.9)	NS
Plurity, n (%)							
Singleton	74 (66.7)	14 (63.6)	60 (67.4)	NS	2 (40.0)	12 (70.6)	NS
Twin/triplet	37 (33.3)	8 (36.4)	29 (32.6)		3 (60.0)	5 (29.4)	
Monochorionicity, n (%)							
Total	9 (24.3)	2 (25.0)	7 (24.1)	NS	1 (33.3)	1 (20.0)	NS
TTTS	6 (66.7)	2 (100)	4 (57.1)	NS	1 (100)	1 (100)	NS
Donor/recipient	4/2	1/1	3/1	NS	0/1	1/0	NS
Congenital TORCH, n (%)							
Performed	24 (27.9)	10 (45.5)	14 (20.9)	NS	2 (40.0)	8 (88.9)	NS
Positive	0 (0)	0 (0)	0 (0)	NS	0 (0)	0 (0)	NS
Sepsis/NEC, n (%)	51 (45.9)	11 (50.0)	40 (44.9)	NS	2 (40.0)	9 (52.9)	NS
RDS, n (%)	61 (55.0)	15 (68.2)	46 (51.7)	NS	4 (80.0)	11 (64.7)	NS
BPD, n (%)	5 (4.5)	1 (4.5)	4 (4.5)	NS	1 (20.0)	0 (0)	NS
Postnatal dexamethasone, n (%)	16 (14.4)	2 (9.1)	14 (15.7)	NS	0 (0)	2 (11.8)	NS
Hypotension, n (%)	37 (33.3)	3 (13.6)	34 (38.2)	0.042*	1 (20.0)	2 (11.8)	NS
PDA, n (%)	36 (32.4)	5 (22.7)	31 (34.8)	NS	1 (20.0)	4 (23.5)	NS

Table 2 Characteristics of clinical parameters for the total group, the lenticulostriate vasculopathy (LSV) group, the non-LSV group, and the earlyonset and late-onset LSV groups, and comparison of clinical parameters between groups

*Independent variable.

BPD, bronchopulmonary dysplasia; GA, gestational age; NEC, necrotising enterocolitis; NS, not significant; PDA, patent ductus arteriosus; RDS, respiratory distress syndrome; TORCH, Toxoplasmosis, Other, Rubella, Cytomegalovirus, Herpes (including Parvo B19); TTTS, twin-to-twin transfusion syndrome.

incidence of 4.6% was reported.¹¹ Possible explanations for our higher incidence may be differences in study populations and ultrasound techniques. We included only very preterm infants, assessed with sequential cUS over a longer period, up to TEA, and also included infants with a punctate appearance of LSV. Ultrasound equipment and techniques have improved considerably over recent years, resulting in higher image quality and possibly better detection of subtle changes. In a study by Paczko et al¹⁰ in a group of preterm infants up to 37 weeks' GA with otherwise normal cUS in the first postnatal week, an incidence of 31.6% was described. Higher incidences have been reported in full-term than in preterm neonates, probably explaining their high incidence. As we included a large cohort of unselected infants born very prematurely, assessed with frequent highquality cUS, our data probably reflect the true incidence of LSV in very preterm infants.

LSV was mostly detected after the first few weeks of birth, and persisted for several months, at least up to TEA. At first detection, LSV was mostly unilateral (mainly right-sided) and linear, but often became bilateral and more complex in appearance on subsequent scans before fading towards TEA. These findings are largely consistent with recent studies, describing a mean postnatal age at first detection of 4–5 weeks, and LSV to be mostly unilateral (right-sided) and have a progressive or static appearance.^{4–9 12} ¹³ We and others do not have an explanation for the right-sided predominance of LSV.

In agreement with previous studies,^{4 9 11}¹⁶ the majority (77%) of our preterm infants had late-onset LSV. Only in studies mainly consisting of full-term neonates, LSV was generally present within the first week of life.^{7 12} ¹³ These findings further indicate that in preterm infants LSV mostly presents after the first postnatal week(s).

Similar to Hemachandra *et al*,¹¹ we found infants with lateonset LSV to be younger at birth than those with early-onset LSV. However, despite the differences in GA and postnatal age at first detection of LSV between the early- and late-onset groups, the PMA at first detection was not significantly different. In most infants, LSV was first seen between 30 and 31 weeks' PMA. This may indicate that PMA is an important factor in the development of LSV, rather than GA at birth and postnatal age at first detection, and that the lenticulostriate vessels are most vulnerable or prone for LSV before or around this PMA. This hypothesis is supported by the fact that in full-term neonates LSV mostly presents at birth, while in preterm neonates mostly after several weeks.^{4 7 9 11–13 16}

LSV has been associated with a variety of perinatal conditions. We found a tendency towards more multiple pregnancies in the early-onset group, being in agreement with several previous studies, reporting LSV to occur more often in infants of multiple pregnancies, particularly monochorionic twin pregnancies.^{12 13 25 26} As we did not find differences in incidence of monochorionic twins between infants with and without LSV, we cannot confirm this finding.

We did not find an association between LSV and congenital TORCH infection: in all infants, both with and without LSV, in whom TORCH screening was performed, results were negative. Our findings are consistent with recent studies, only occasionally describing TORCH infection in neonates with LSV.^{7 11 13} Therefore, we do not consider TORCH screening warranted in very preterm infants with LSV, unless there are symptoms suggestive of infection.

The aetiology and clinical significance of LSV remain largely unclear.⁸ ¹³ ¹⁵ While initial studies suggested an infectious origin,^{1–} ³ ¹⁹ more recent studies have suggested (a combination of) a variety of infectious and non-infectious origins.⁶⁷⁹ In addition, late-onset LSV may be due to a perinatal insult to the developing brain, while early-onset LSV may reflect an in utero insult.¹¹

Histological studies on changes in thalamo-striatal vessels in infants with LSV reported inconsistent findings, possibly partly due to sampling differences and to different stages of LSV.⁵ ⁹ Some studies found no (peri)vascular abnormalities in involved vessel walls, while others reported depositions of basophilic

Original article

material as a solitary finding or in combination with iron (mineralisation), diffuse microcalcifications and/or calcium.^{2 3 5 9 13} LSV was therefore thought to represent a vasculopathy, but its underlying pathogenesis is unclear and several potential pathways have been suggested, including vasculitis, hypoxia–ischaemia, and changes in blood flow.

We only found a significant association between LSV, particularly late-onset, and less episodes of hypotension, and do not have an explanation for this finding. We did not find associations with (other) changes on cUS or MRI, and, as expected, no equivalent was found for LSV on MRI.²⁻⁶ 13 21 22 28 An explanation for this may be that ultrasound and MR imaging are based on different technologies, making detection of vascular deposits and/or calcifications difficult with MRI. In addition, LSV seems to be of vascular origin and does not represent tissue changes. Therefore, the MRI techniques included in this study may not be sensitive to the changes underlying LSV. Possibly, studies on blood flow in and vasculature of involved vessels, including MR angiography, may elucidate vascular changes resulting from LSV and/or may help to study its origin. Based on our results and recent findings, we suggest that a combination of factors underlies LSV.

We appreciate some limitations of our study. In several infants early transfer limited sequential cUS to the early postnatal period. As in most infants LSV was still seen at discharge, it was not always possible to assess its exact duration, and we may have underestimated its duration in four infants in whom it was no longer detected around TEA. We excluded infants with congenital malformations and chromosomal and/ or metabolic disorders. Consequently, we cannot comment on the association between LSV and these disorders, and may have slightly underestimated the incidence of LSV in the general preterm population. Finally, no outcome data are, as yet, available for our study population. Favourable outcome has been found in infants with isolated LSV.¹³

In conclusion, this study shows that LSV is a frequent finding in very preterm infants. It often first presents several weeks after birth and persists for several months. PMA, rather than GA and postnatal age, is an important factor in the development of LSV. The aetiology and clinical significance remain unclear and we did not find associations with perinatal clinical parameters, including congenital infections. Our findings suggest that LSV, when encountered in otherwise healthy preterm infants, may be a benign temporary phenomenon.

Acknowledgements: We are grateful to the staff of the neonatal unit, the paediatric ambulatory unit and the MRI department of the Leiden University Medical Center and to B. van Kooij, University Medical Center Utrecht, for their helpful contributions to this project. We thank W. Teeuwisse for technical support.

Funding: This project was supported by a grant 920-03-388 from ZonMw, the Netherlands Organisation for Health Research and Development.

Competing interests: None.

Ethics approval: The study was approved by the Medical Ethics Committee.

Patient consent: Parental consent obtained.

Provenance and peer review: Not commissioned; externally peer reviewed.

REFERENCES

- Grant EG, Williams AL, Schellinger D, et al. Intracranial calcification in the infant and neonate: evaluation by sonography and CT. Radiology 1985;157:63–8.
- Teele RL, Hernanz-Schulman M, Sotrel A. Echogenic vasculature in the basal ganglia of neonates: a sonographic sign of vasculopathy. *Radiology* 1988;169:423–7.
- Hughes P, Weinberger E, Shaw DWW. Linear areas of echogenicity in the thalami and basal ganglia of neonates: an expanded association. Work in progress. *Radiology* 1991;179:103–5.
- Weber K, Riebel Th, Nasir R. Hyperechoic lesions in the basal ganglia: an incidental sonographic finding in neonates and infants. *Pediatr Radiol* 1992;22:182–6.

- Cabañas F, Pellicer A, Morales C, et al. New pattern of hyperechogenicity in thalamus and basal ganglia studied by color Doppler flow imaging. *Pediatr Neurol* 1994;10:109–16.
- Wang HS, Kuo MF, Chang TC. Sonographic lenticulostriate vasculopathy in infants: some associations and a hypothesis. *AJNR Am J Neuroradiol* 1995;16:97–102.
- Shefer-Kaufman N, Mimouni FB, Stavorovsky Z, et al. Incidence and clinical significance of echogenic vasculature in the basal ganglia of newborns. Am J Perinatol 1999;16:315–9.
- Chamnanvanakij S, Rogers CG, Luppino C, *et al*. Linear hyperechogenicity within the basal ganglia and thalamus of preterm infants. *Pediatr Neurol* 2000;23:129–33.
 Coley BD, Rusin JA, Boue DR. Importance of hypoxic/ischemic conditions in the
- development of cerebral lenticulostriate vasculopathy. *Pediatr Radiol* 2000;**30**:846–55.
 Paczko N, Rotta NT, Silva A, *et al.* Hyperechogenicity of thalamic vessels in preterm
- newborn infants. J Pediatr (Rio J) 2002;78:371–4.
 Hemachandra AH, Oravec D, Collin M, et al. Early and late postnatal identification of isolated lenticulostriate vasculopathy in preterm infants: associated findings. Perinatol 2003:23:20–3.
- Makhoul IR, Eisenstein I, Sujov P, et al. Neonatal lenticulostriate vasculopathy: further characterisation. Arch Dis Child Fetal Neonatal Ed 2003;88:F410–4.
- El Ayoubi M, de Bethmann O, Monset-Couchard M. Lenticulostriate echogenic vessels: clinical and sonographic study of 70 neonatal cases. *Pediatr Radiol* 2003;33:697–703.
- Mittendorf R, Covert R, Pryde PG, et al. Association between lenticulostriate vasculopathy (LSV) and neonatal intraventricular hemorrhage (IVH). J Perinatol 2004;24:700–5.
- Mittendorf R, Kuban K, Pryde PG, et al. Antenatal risk factors associated with the development of lenticulostriate vasculopathy (LSV) in neonates. J Perinatol 2005;25:101–7.
- Shen EY, Weng SM, Kuo YT, et al. Serial sonographic findings of lenticulostriate vasculopathy. Acta Paediatr Taiwan 2005;46:77–81.
- Cabañas F, Pellicer A, Valverde E, *et al.* Central nervous system vasculopathy in neonatal lupus erythematosus. *Pediatr Neurol* 1996;15:124–6.
- Tomà P, Magnano GM, Mezzano P, et al. Cerebral ultrasound images in prenatal cytomegalovirus infection. *Neuroradiology* 1989;31:278–9.
- Ben-Ami T, Yousefzadeh D, Backus M, et al. Lenticulostriate vasculopathy in infants with infections of the central nervous system: sonographic and Doppler findings. Pediatr Radiol 1990;20:575–9.
- Yamashita Y, Matsuishi T, Murakami Y, et al. Neuroimaging findings (ultrasonography, CT, MRI) in 3 infants with congenital rubella syndrome. *Pediatr Radiol* 1991;21:547–9.
- de Vries LS, Gunardi H, Barth PG, et al. The spectrum of cranial ultrasound and magnetic resonance imaging abnormalities in congenital cytomegalovirus infection. *Neuropediatrics* 2004;35:113–9.
- Chabra S, Kriss VM, Pauly TH, et al. Neurosonographic diagnosis of thalamic/basal ganglia vasculopathy in trisomy 13: an important diagnostic aid. Am J Med Genet 1997;72:291–3.
- Lin HY, Lin SP, Chen YJ, et al. Clinical characteristics and survival of trisomy 13 in a medical center in Taiwan, 1985–2004. *Pediatr Int* 2007;49:380–6.
- te Pas AB, van Wezel-Meijler G, Bökenkamp-Gramann R, et al. Preoperative cranial ultrasound findings in infants with major congenital heart disease. Acta Paediatr 2005;94:1597–603.
- de Vries LS, Beek FJA, Stoutenbeek P. Lenticulostriate vasculopathy in twin-to-twin transfusion syndrome: sonographic and CT findings. *Pediatr Radiol* 1995;25:S41–2.
- Kandasamy Y, Alcock G, Koh THHG. Lenticulostriate vasculopathy in twin-to-twin transfusion syndrome. J Perinatol 2006;26:780–2.
- Lopriore E, van Wezel-Meijler G, Middeldorp JM, et al. Incidence, origin, and character of cerebral injury in twin-to-twin transfusion syndrome treated with fetoscopic laser surgery. Am J Obstet Gynecol 2006;194:1215–20.
- Leijser LM, de Vries LS, Rutherford MA, et al. Cranial ultrasound in metabolic disorders presenting in the neonatal period: characteristic features and comparison with MR imaging. AJNR Am J Neuroradiol 2007;28:1223–31.
- van Wezel-Meijler G. Neonatal cranial ultrasonography. Heidelberg: Springer Verlag, 2007.
- Leijser LM, de Bruïne FT, Steggerda SJ, et al. Brain imaging findings in very preterm infants throughout the neonatal period: Part I. Incidences and evolution of lesions, comparison between ultrasound and MRI. Early Hum Dev 2009;85:101–9.
- van Wezel-Meijler G, Leijser LM, de Bruïne FT, et al. Magnetic resonance imaging of the brain in newborn infants: practical aspects. *Early Hum Dev* 2009;85:85–92.
- Barkovich AJ. Normal development of the neonatal and infant brain, skull and spine. In: Barkovich AJ, ed. *Pediatric neuroimaging*. Philadelphia: Lippincott Williams & Wilkins, 2005: 13.
- Battin M, Rutherford MA. Magnetic resonance imaging of the brain in preterm infants: 24 weeks' gestation to term. In: Rutherford MA, ed. *MRI of the neonatal brain*. Edinburgh: W.B. Saunders, 2002: 25–50.
- Cowan FM. Magnetic resonance imaging of the normal infant brain: term to 2 years. In: Rutherford MA, ed. *MRI of the neonatal brain*. Edinburgh: W.B. Saunders, 2002: 51–84.
- van der Knaap MS, Valk J. Myelination and retarded myelination. In: van der Knaap MS, Valk J, eds. *Magnetic resonance of myelination and myelin disorders*. Berlin: Springer Verlag, 2005: 37–65.
- Bancalari E, Claure N. Definitions and diagnostic criteria for bronchopulmonary dysplasia. Semin Perinatol 2006;30:164–70.
- Bell MJ, Ternberg JL, Feigin RD, et al. Neonatal necrotizing enterocolitis. Therapeutic decisions based upon clinical staging. Ann Surg 1978;187:1–7.